

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Comparison of different procedures for the determination of element bioaccessibility in ayurvedic formulationss

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1695870> since 2019-03-30T21:10:38Z

Publisher:

XVI Hungarian – Italian Symposium on Spectrochemistry

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

COMPARISON OF DIFFERENT PROCEDURES FOR THE DETERMINATION OF ELEMENT BIOACCESSIBILITY IN AYURVEDIC FORMULATIONS

**Ornella Abollino¹, Agnese Giacomino², Eleonora Conca¹, Valentina Barbero²,
Diana Toma², Sandro Buoso¹, Marina Di Carro³, Mery Malandrino¹**

¹*Dipartimento di Chimica, Università di Torino,
Via Giuria 7, 10125 Torino, Italy*

²*Dipartimento di Scienza e Tecnologia del Farmaco, Università di Torino,
Via Giuria 9, 10125 Torino, Italy*

³*Dipartimento di Chimica e Chimica Industriale, Università di Genova,
Via Dodecaneso 31, 16146 Genova
e-mail: ornella.abollino@unito.it*

The knowledge of the bioavailability and bioaccessibility of elements in foods, food supplements and medicines is very important to assess the potential health hazard associated to their consumption, since not all the amount of an element present in an ingested product is available for absorption by the organism. Oral bioaccessibility is the fraction of a compound that is released from its matrix in the gastrointestinal tract and thus becomes available for intestinal absorption [1]. It is determined *in vitro* by extraction with solutions simulating gastric and intestinal fluids and provides a precautionary estimation of bioavailability, since the extracted substance might not be completely absorbed. In this work we have determined the total and bioaccessible concentrations of Al, As, Ba, Cd, Cu, Fe, Hg, Ni, Pb and Zn in a suite of ayurvedic products belonging to the *rasausadhi* tradition, which relies on the deliberate combination of herbs, metals and minerals; the products were purchased in India and are not available in the UE, where only metal-free ayurvedic products are admitted. The work is the prosecution of a previous study dealing with other Ayurvedic products [2]. In the present research we have put the samples in contact with the extractants in different ways, to investigate the effect of experimental conditions on extractability: i) the samples were simply poured into the liquid; ii) they were mixed with *ghee*, the Indian butter, according to the traditional way of assumption of ayurvedic products; iii) they were suspended in the liquid with the aid of a semipermeable membrane. Concentrations were determined by Inductively Coupled Plasma-Optical Emission Spectroscopy (ICP-OES) after microwave digestion. The results showed that bioaccessibility decreased using the second and third procedure. Bioaccessible concentrations were lower than total concentrations and extractability was higher in synthetic gastric fluids than in intestinal ones.

References

- [1] M. Intawongse, J.R. Dean, In-vitro testing for assessing oral bioaccessibility of trace metals in soil and food samples, *TrAC – Trend Anal Chem.* 25 (2006) 876–86.
- [2] A. Giacomino, O. Abollino, C. Casanova, C. La Gioia, E. Magi, M. Malandrino, Determination of the total and bioaccessible contents of essential and potentially toxic elements in ayurvedic formulations purchased from different commercial channels, *Microchem. J.* 120 (2015) 6–17.